

Product Information: Similac® Advance®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Advance®

Infant Formula with Iron

- 1 A nutritionally complete, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding.
- 1 Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
- 1 Clinically proven tolerance in newborns and proven immune outcomes.
- 1 Similac Advance has been clinically shown to support greater calcium absorption* for strong bones.
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* *Similac® with Iron vs Enfamil® with Iron. Enfamil is not a trademark of Abbott Laboratories.*

Ingredients

Powder Unflavored:

Nonfat Milk, Lactose, Whey Protein Concentrate, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, m-Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Calcium Phosphate, Potassium Phosphate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Availability

List Number	Item
53359	Similac Advance Powder / 1.45-lb (658-g) SimplePac / 6 ct
53363	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 6 ct
53365	Similac Advance Ready to Feed Institutional / 2-fl-oz (59-mL) bottle / 48 ct
53496	Similac Advance Powder / 2.13-lb (964-g) SimplePac / 6 ct
53498	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 6 ct
53526	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 12 x 1 ct
55957	Similac Advance Powder / 12.4-oz (352-g) can / 6 ct
56168	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 8 ct
56973	Similac Advance Concentrated Liquid / 13-fl-oz (384-mL) can / 12 ct
57600	Similac Advance Newborn Bottles Ready to Feed / 2-fl-oz (59 mL) bottle / 6 x 8 ct
57938	Similac Advance On-the-go Powder Singles / 0.61-oz (17.4-g) packet / 6 x 16 ct
58585	Similac Advance Powder / 1.93-lb (873-g) can / 6 ct
58605	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 4 x 6 ct
59952	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 24 ct
62811	Similac Advance On-the-go Powder Singles / 0.61-oz (17.4-g) packet / 4 x 16 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.07	14.00
Fat, g	5.60	37.8
Carbohydrate, g	10.71	72.4
Water, g	133	899
Linoleic Acid, mg	1000	6757
Potential Renal Solute Load, mOsm	18.7	126.7
Vitamin A, IU	300	2029
Vitamin D, IU	75	507
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.25	1.69
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.7	31.8
Calcium, mg	78	528
Calcium, mEq	3.9	26.3
Phosphorus, mg	42	284
Magnesium, mg	6	41
Iron, mg	1.8	12
Zinc, mg	0.75	5.07
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	162
Sodium, mEq	1.04	7.1
Potassium, mg	105	710
Potassium, mEq	2.7	18.2
Chloride, mg	65	440
Chloride, mEq	1.8	12.4

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.

0.61-oz powder packet

- 1 Pour **4 fl oz** of water into clean bottle.
- 1 Slowly add one powder single (17.4 g) to bottle.
- 1 Cap bottle; shake very well; attach nipple and ring.
- 1 Once feeding begins, **use within 1 hour or discard**.

12.4-oz can, 1.45-LB, 1.93-LB & 2.13-LB SimplePac containers

- 1 Pour water into clean bottle (see mixing guide).
- 1 Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- 1 Return dry scoop to container.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, **use within 1 hour or discard**.

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Concentrated Liquid

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

13-fl-oz can

- 1 Rinse can lid; shake very well; open with a clean punch-type opener.
- 1 Pour equal amounts of formula and water into bottle.
- 1 Attach nipple; swirl gently to mix.
- 1 Once feeding begins, **use within 1 hour or discard**.

1-QT bottle

- 1 Shake very well before each use.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

- 1 Remove protective seal around bottle cap; twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour equal amounts of formula and water into bottle.
- 1 Attach nipple; swirl gently to mix.
- 1 Once feeding begins, **use within 1 hour or discard.**
- 1 **Prepare and feed all formula from this container within 4 days after opening.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Do not add water.
- 1 Shake very well.
- 1 Twist off cap.
- 1 Twist on nipple and ring.
- 1 No need to warm before feeding.
- 1 Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Do not add water.
- 1 Shake very well.
- 1 Twist off cap.
- 1 Twist on clean nipple and ring.
- 1 No need to warm before feeding.
- 1 Once feeding begins, **use within 1 hour or discard.**
- 1 For use in a different feeding container (cup or bottle):
 - 1 Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - 1 Pour desired amount into clean feeding container.
 - 1 Recap bottle, refrigerate and **use within 48 hours.**

1-QT bottle

- 1 Do not add water.
- 1 Shake very well before each use.
- 1 Remove protective band, twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle; attach nipple.
- 1 Once feeding begins, **use within 1 hour or discard**
- 1 Prepare and feed all formula from the 1-QT container **within 3 days after opening.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Do not add water.
- 1 Shake very well.
- 1 Twist off cap.
- 1 Twist on nipple and ring.
- 1 No need to warm before feeding.
- 1 Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Do not add water.
- 1 Shake very well.
- 1 Twist off cap.
- 1 Twist on clean nipple and ring.
- 1 No need to warm before feeding.
- 1 Once feeding begins, **use within 1 hour or discard.**
- 1 For use in a different feeding container (cup or bottle):
 - 1 Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - 1 Pour desired amount into clean feeding container.
 - 1 Recap bottle, refrigerate and **use within 48 hours.**

1-QT bottle

- 1 Do not add water.
- 1 Shake very well before each use.
- 1 Remove protective band, twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle; attach nipple.
- 1 Once feeding begins, **use within 1 hour or discard**
- 1 Prepare and feed all formula from the 1-QT container **within 3 days after opening.**

Storage & Handling

Powder

0.61-oz powder packet

- 1 Once mixed, feed immediately or cover and refrigerate; **use within 24 hours.**
- 1 Store unopened packets at room temperature; avoid extreme temperatures.

12.4-oz can, 1.45-LB, 1.93-LB & 2.13-LB SimplePac containers

- 1 Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- 1 Store unopened or opened container at room temperature; avoid extreme temperatures.
- 1 **Use opened can contents within 1 month.**
- 1 Do not reuse container.

Concentrated Liquid

13-fl-oz can

- 1 Cover opened cans and prepared formula not used immediately. Store in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened cans at room temperature; avoid extreme temperatures.

1-QT bottle

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened container at room temperature; avoid extreme temperatures.
- 1 Do not reuse container.

Ready to Feed

2-fl-oz bottle & 8-fl-oz bottle

- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Do not reuse bottle.
- 1 **Protect from light.**
- 1 **Store in carton until ready to use.**

1-QT bottle

- 1 Once opened, store quart bottle in refrigerator
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Do not reuse container.